


Claims

1. **(original)** A time temperature indicator for indicating a temperature change over time, comprising at least one indicator compound in a first isomeric form, which is converted into second isomeric form of said indicator compound in a valence isomerization reaction without migration of an atom or chemical group attached to said indicator compound in a time and temperature dependent manner, wherein the formation of the second isomeric form is detectable by monitoring a physical characteristic of the first isomeric or the second isomeric form of the indicator.
2. **(original)** The time-temperature indicator of claim 1, wherein the at least one indicator compound is a diarylethene or a spiroaromatic compound.
3. **(currently amended)** The time-temperature indicator **[[of]]** of claim 2, wherein the diarylethene is a compound of Formula (I)



wherein

R1 and R2 each independently represent C6-C14 aryl, C4-C12 heteroaryl, conjugated heterocyclic; wherein said heteroaryl and conjugated heterocyclic may contain one to three heteroatoms selected from N, O, **[[or]]** **and** S; and wherein said aryl, heteroaryl, or conjugated heterocyclic may be substituted by one or more halogen, hydroxyl, thiol, amino, C1-C12 alkyl, C2-C12 alkenyl, C2-C12 alkynyl, C1-C6 alkanoyl, C1-C6 alkoxy, C1-C6 alkylthio, C6-C14 aryl, C4-C14 heteroaryl, C3-C8 membered non-aromatic carbocyclic, C3-C8 membered ring non-aromatic heterocyclic, cyano, nitro, sulfo, -CH=CH-CN, azido, or amido;

R1' and R2' each independently represent H, cyano, nitro, sulfo, hydroxyl, thiol, -CH=CH-CN, or amido; or substituted or unsubstituted C1-C12 alkyl, C2-C12 alkenyl, C2-C12 alkynyl, C1-C6 alkanoyl, C1-C6 alkoxy, C1-C6 alkylthio, C6-C14 aryl, C4-C14 heteroaryl, C3-C8 membered non-aromatic carbocyclic, C3-C8 membered ring non-aromatic heterocyclic; or R1' and R2' together with the carbon atoms to which they are attached form a C5-C8 carbocyclic ring or a


C4-C7 heterocyclic ring containing one to three endocyclic or exocyclic heteroatoms selected from N, O, **[[or]] and** S; said N heteroatom may be further substituted by H, or by one or two substituted or unsubstituted groups selected from C1-C12 alkyl, C2-C12 alkenyl, C2-C12 alkynyl, C1-C6 alkanoyl, C1-C6 alkoxy, C1-C6 alkylthio, C6-C14 aryl, C4-C14 heteroaryl, C3-C8 membered non-aromatic carbocyclic, C3-C8 membered ring non-aromatic heterocyclic, hydroxyl, **[[or]] and** -CH=CH-CN; when said N heteroatom is tetrasubstituted it is positively charged and is associated with an anion selected from the group consisting of organic **[[or]] and** inorganic anions, and optionally wherein said C5-C8 carbocycle is substituted by one or more halogen, preferably by one or more fluoro atoms; and optionally

R1, R1', R2 and R2' each independently represent a charged group or a group substituted by another group having a charge; said charge may be localized or delocalized and may be positive or negative;

and wherein said R1 and R2 are in a cis or trans conformation.

4. **(currently amended)** The time-temperature indicator of claim 3, wherein the diarylethene is
- (a) a symmetric diarylethene, ~~preferably~~ selected from the group consisting of 1,2-dicyano-1,2-bis(2,4,5-trimethylthiophene-3-yl)ethane (1); 2,3-bis(2,4,5-trimethylthiophene-3-yl)maleic anhydride (2); 1,2-bis(2-cyano-1,5-dimethyl-4-pyrrolyl)perfluorocyclopentene (3); and 1,2-bis(2,4-dimethyl-5-phenylthiophene-3-yl)perfluorocyclopentene (4); or
 - (b) an asymmetric diarylethene, ~~preferably~~ selected from the group consisting of 2-(1,2-dimethyl-3-indolyl)-3-(2,4,5-trimethyl-3-thienyl) maleic anhydride (5); and 2-(methoxybenzo[b]thiophene-3-yl)-3-(1,2-dimethyl-3-indolyl) maleic anhydride (6).

5. **(currently amended)** The time-temperature indicator of claim 2, wherein the spiroaromatic compound is a compound of Formula (II):



wherein

ring A represents a C5-C8 carbocycle, C4-C7 heterocycle containing at least one heteroatom selected from N, O, **[[or]] and** S; said N heteroatom may be further substituted by one or two

groups selected from C1-C12 alkyl, C2-C12 alkenyl, C2-C12 alkynyl, C1-C6 alkanoyl, C1-C6 alkoxy, C1-C6 alkylthio, C6-C14 aryl, C4-C14 heteroaryl, C3-C8 membered non-aromatic carbocyclic, C3-C8 membered ring non-aromatic heterocyclic, hydroxyl, ~~[[or]]~~ **and** -CH=CH-CN; when said N heteroatom is tetrasubstituted it is positively charged and is associated with an anion selected from the group consisting of organic or inorganic anions; said C5-C8 carbocycle or C4-C7 heterocycle may be substituted by one or more of the groups selected from halogen, C1-C12 alkyl, C2-C12 alkenyl, C2-C12 alkynyl, C1-C6 alkanoyl, C1-C6 alkoxy, C1-C6 alkylthio, C6-C14 aryl, C4-C14 heteroaryl, C3-C8 membered non-aromatic carbocyclic, C3-C8 membered ring non-aromatic heterocyclic, cyano, nitro, sulfo, hydroxyl, thiol, -CH=CH-CN, azido, amido ~~[[or]]~~ **and** amino;

ring B represents a substituted or unsubstituted heterocycle containing at least one heteroatom X, said X being selected from N, O, and S; wherein said N atom may be further substituted by one or two groups selected from C1-C12 alkyl, C2-C12 alkenyl, C2-C12 alkynyl, C1-C6 alkanoyl, C1-C6 alkoxy, C1-C6 alkylthio, C6-C14 aryl, C4-C14 heteroaryl, C3-C8 membered non-aromatic carbocyclic, C3-C8 membered ring non-aromatic heterocyclic, hydroxyl, or -CH=CH-CN; when said N heteroatom is tetrasubstituted it is positively charged and is associated with an anion selected from the group consisting of organic or inorganic anions;


and wherein said ring B may contain one or more endocyclic double bonds and is optionally substituted by one or more halogen, ~~preferably by one or more fluoro atoms;~~

said rings A and B may be fused to one or more substituted or unsubstituted carbocycle, C4-C14 heterocycle, C6-C14 aryl or C4-C14 heteroaryl ring system;

and wherein the compounds of Formula II may be neutral, charged, multiply charged, positively charged having an external anion, negatively charged having an external cation or zwitterionic.

6. **(currently amended)** The time-temperature indicator of claim 5, wherein the spiroaromatic compound is a spiropyran derivative, ~~preferably selected from the group consisting of 1',3',3',8-tetramethyl-5-hydroxymethyl-spiro(2H-pyrano[2,3-c]pyridine-2,2'-2H-indole) and 1',3',3',8-tetramethyl-spiro(2H-pyrano[2,3-c]pyridine-2,2'-2H-indole).~~

7. **(currently amended)** The time-temperature indicator of claim 5, wherein the spiropyran derivative is a derivative of 1',3',3'-trimethyl-6-nitro-spiro(2H-1-benzopyran-2,2'-2H-indole) of Formula (III):



wherein


R3 is selected from the group consisting of H, halogen, C1-C12 alkyl, C2-C12 alkenyl, C2-C12 alkynyl, C1-C6 alkanoyl, C1-C6 alkoxy, C1-C6 alkylthio, C6-C14 aryl, C4-C14 heteroaryl, C3-C8 membered non-aromatic carbocyclic, C3-C8 membered ring non-aromatic heterocyclic, **[[or]]** **and** azido; wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, and non-aromatic carbocycle may be substituted by one or more group selected from halogen, hydroxyl, thiol, amino, alkoxy, nitro, azido, **[[or]]** **and** sulfo;

R4 is selected from the group consisting of C1-C12 alkyl, C2-C12 alkenyl, C2-C12 alkynyl, C1-C6 alkanoyl, C1-C6 alkoxy, C1-C6 alkylthio, C6-C14 aryl, C4-C14 heteroaryl, C3-C8 membered non-aromatic carbocyclic, C3-C8 membered ring non-aromatic heterocyclic, hydroxyl **[[or]]** **and** -CH=CH-CN; and

Y is selected from the group consisting of C1-C25 alkyl, ~~preferably methyl, n-propyl and n-octadecyl,~~ and C7-C15 aralkyl, wherein said alkyl and aralkyl ~~may be~~ is optionally substituted by one or more ~~group selected from halogen, preferably fluorine.~~

8. **(currently amended)** The time-temperature indicator of **[[C]]** claim 5, wherein the spiroaromatic compounds include at least one of the following: spirooxazine or its derivatives, spironaphthoxazine or its derivatives, and spiroindolinopyridobenzoxazine or its derivatives.

9. **(currently amended)** A spiroaromatic compound of general Formula (IV):



wherein

A and L are independently of each other selected from the group consisting of H, halogen, C2-

C12 alkenyl, C2-C12 alkynyl and $\text{C}(=\text{O})\text{R}$, wherein R is C1-C6 alkyl, C1-C6 alkoxy, C6-C14 aryl

and C7-C15 aralkyl; wherein said alkenyl, alkynyl and $\text{C}(=\text{O})\text{R}$ may be substituted by one or more group selected from halogen, hydroxyl, thiol, amino, alkoxy, nitro, azido, sulfo, aryl and heteroaryl;

Y is selected from the group consisting of C1-C25 alkyl and C7-C15 aralkyl, wherein said alkyl and aralkyl ~~may be~~ is optionally substituted by one or more ~~group selected from halogen,~~ preferably fluorine; and

X is C1-C6 alkoxy or L;

with the proviso that Y is not n-propyl when L, A and X are hydrogen.

10. **(original)** The spiroaromatic compound of claim 9, wherein

L is hydrogen, Cl, Br or I;

Y is methyl, n-propyl, n-octadecyl or $\text{F}-\text{C}_6\text{H}_4-\text{CH}_2-$;

X is hydrogen or methoxy; and

A is hydrogen;

with the proviso that Y is not n-propyl when L and X are hydrogen.

11. **(currently amended)** A printing ink or printing ink concentrate, comprising the spiroaromatic compound of claim 9, ~~or 10.~~

12. **(currently amended)** A high molecular weight material, comprising the spiroaromatic compound of claim 9, ~~or 10.~~

13. **(currently amended)** A method of manufacturing a time-temperature indicator of claim 1 ~~any of claims 1 to 8~~ comprising the steps of
- (a) embedding in or atop a matrix said at least one indicator compound; and
 - (b) inducing the formation of a metastable state of said embedded at least one indicator compound.
14. **(currently amended)** The method of claim 13 ~~[[9]]~~, further comprising the step of covering the time-temperature indicator with a cover support, ~~preferably designed to avoid photo recharging and/or photo bleaching.~~
15. **(new)** The time-temperature indicator of claim 6, wherein the spiropyran derivative is selected from the group consisting of 1',3',3',8-tetramethyl-5-hydroxymethyl-spiro(2H-pyrano[2,3-c]pyridine-2,2'-2H-indole) and 1',3',3',8-tetramethyl-spiro(2H-pyrano[2,3-c]pyridine-2,2'-2H-indole).
16. **(new)** The time-temperature indicator of claim 7, wherein in Formula (III) Y is selected from the group consisting of C1-C25 alkyl and C7-C15 aralkyl, wherein said alkyl and aralkyl are substituted by one or more fluorine.
17. **(new)** A spiroaromatic compound of claim 9, wherein in Formula (IV) Y is selected from the group consisting of C1-C25 alkyl and C7-C15 aralkyl, wherein said alkyl and aralkyl are substituted by one or more fluorine.
18. **(new)** A printing ink or printing ink concentrate, comprising the spiroaromatic compound of claim 10.
19. **(new)** A high molecular weight material, comprising the spiroaromatic compound of claim 10.
20. **(new)** The method of claim 14, wherein the cover support is designed to avoid photo recharging and/or photo bleaching.